



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

- I. *A Catalogue of the fifty Plants from Chelsea-Garden, presented to the Royal Society by the Company of Apothecaries, for the Year 1730; pursuant to the Direction of Sir Hans Sloane, Bart. Med. Reg. Pres. Col. Reg. Med. & Soc. Reg. by Isaac Rand, Apothecary, F. R. S.*

401. **A** DHATODA Zeylanensium. Hort. Lugd. Bat. 642.

402. Ageratum foliis ferratis. C. B. 221. *Ageratum plerisque; Herba Julia quibusdam J. B.* III. pag. 142.

403. Ageratum, breviori & latiori folio. *Ptarmitica lutea, suaveolens; corymbis longioribus & magis compactis.* T. Inst. 497.

404. Ageratum Hispanicum; amplioribus foliis; flosculis minoribus.

405. Anonis pulchra visu; folio Ciceris; Ornithopodij filiquis, annuentibus; Hort. Cath. 18.

406. Aparine femine Coriandri saccharati. Park. Theat. 567.

407. Aparine femine læviore, Raij Hist. 484.

408. Aparine minima, D. Sherard. Raij Syn. Ed. 2. 118.

409. Arundo arborea Indica; amplo, pediculato folio. *Bambu altera species.* Raij Hist. 1316. *Beesha.* Hort. Malab. Tom. 5. Tab. 60.

410. *Canna Indica* ; radice alba, alexipharmica, Sloan. Cat. Jam. 122. Hist. Tab. 149. Fig. 2. *Arrow-root*.

411. *Caryophyllata vulgaris*. C. B. 321.

412. *Caryophyllata, aquatica, nutante flore* C. B. 321.

413. *Caryophyllata major* ; nutante flore.

414. *Caryophyllata Virginiana* ; albo flore, minore ; radice inodora. H. Lugd. Bat. 121.

415. *Caucalis Monspeliaca* ; echinato magno fructu. C. B. 153.

416. *Chamædrys Alpina*, frutescens ; folio splendente. T. Inst. 205.

417. *Chamæmelum montanum* ; Absinthij folio ; Parthenij odore. Hort Cath.

418. *Chamæmelum incanum* ; Absinthij vulgaris folio, D. Vaillant. Ac. Reg. Sc. 1720.

419. *Chamæmelum Hispanicum*, humile, ramosissimum ; *Cotulæ foetide* facie.

420. *Chelone Acadiensis* ; flore albo. D. Tournef. Ac. Reg. Sc. 1706. Josselyn *New-England's Rarities*, Numb. 6. pag. 78 & 79.

421. *Eupatoriophalacron*, *Balsaminæ foeminæ* folio ; flore albo discoide. D. Vaillant, Ac. Reg. Sc. 1720. *Scabiosa-Conyzoides Americana* ; capitulis & floribus albidis parvis P. B. P. Pluk. phyt. Tab. 109. Fig. 1.

422. *Eupatoriophalacron Americanum*, procumbens ; albis floribus ; foliis parvis undulatis, asperis.

423. *Eupatoriophalacron Americanum* ; scrophulariæ acutiore folio ; disco luteo, prominente, stipato, vix radiato.

424. Eupatoriophalacron Indicum; scrophulariae folio; capitulo parvo, radiis quinis ad Basin ornato, hispidis & glutinosis.

An Cichoreo affinis Lampfana Sinica, Mentastrifoliis; calyce fimbriato, hispido. Sinice *Hi-him-tsarw dicta?* Pluk. Amalt. 58. Phyt. Tab. 380. Fig. 2.

425. Fœnum Græcum fativum. C. B. 348.

426. Fœnum Græcum, sylvestre; polyceration, majus, Creticum. Breyn. Cent. 79.

427. Fœnum Græcum sylvestre, polyceration, minus, Monspeliacum. Breyn. Cent. 79. *Hedysarum minimum, Dalechamp.* Hist. Lugd. 446.

428. Herniaria glabra. I. B. III. 378.

429. Herniaria hirsuta. I. B. III. 379.

430. Lyfimachia spicata; flore albo; falcis folio. T. Inst. 141.

431. Medica scutellata. I. B. II. 384.

432. Medica hirsuta; echinis rigidioribus. I. B. II. 385.

433. Medica magna, turbinata. I. B. II. 385.

434. Medica tornata, minor, lenis. Park. Theat. 1116.

435. Medica maritima, nostras; modiolis spinosis. *Trifolium cochleatum; modiolis spinosis.* Raij Syn. Ed. 2. 197.

436. Medica fructu ovali spinoso. *Medica διχαρπῶ fructu ovali, Spinulis brevioribus & rigidioribus.* Raij Hist. 963.

437. Medica Orientalis; fructu rotundiore, subhirsuto, tuberculis raris obfuso.

438. Medica Hispanica; fructu echinato, subhirsuto, sphaeroide.

439. *Medica echinata*, minima I. B. II. 386.

440. *Medica marina*. Lob. Icon. 38. *Trifolium cochleatum, maritimum, tomentosum*. C. B. 329.

441. *Melilotus Messanensis*, procumbens; folliculis rugosis, sublongis; spicis florum brevioribus. Raij Hist. 952.

442. *Ornithopodium scorpioides*; siliqua compressa. T. Inst. 400.

443. *Papaver luteum*, perenne; laciniato folio; Cambro-britannicum. Raij Syn. Ed. 3. 309 *Argemone lutea Cambro-britannica*. Park. Theat. 369.

444. *Papaver spinosum*. C. B. 171.

445. *Pseudo-Ipecacuanha Noveboracensis* Dñi Col- den.

An *Periclymenum herbaceum, rectum, Virginianum*, Dr Tinkars-weed, vulgo vocatum? Pluk. Alm. 287. Phyt. Tab. 104. f. 2.

446. *Parmica Agerati folio*, acutiori. *Ageratum album* vulgo.

447. *Ranunculus arvensis*, echinatus. C. B. 179.

448. *Ranunculus arvensis*; seminibus magis echinatis. D. Sherard. Boerh. Ind. alt. 32.

449. *Ruyfchiana*; flore cæruleo, magno. Boerh. Ind. alt. 173. *Chamæpitys cærulea austriaca*. C. B. 250.

450. *Veronica fruticosa erecta*, dulcis; hexangulari caule; flore dilutè cæruleo. Sloan. Cat. Jam. 81. Hist. Tab. 108. f. 2.